



# 2023 LakeForecast End-of-Season Report

## Overview

Clean Lakes Alliance’s LakeForecast water quality monitoring program completed its 11th season in 2023. This program is entirely volunteer-implemented, with 90 trained monitors assessing water quality conditions from nearshore and offshore locations across all five Yahara lakes (Mendota, Monona, Wingra, Waubesa, and Kegonsa). From Memorial Day through Labor Day, volunteers recorded water clarity, air and water temperature, waterfowl presence, extent of floating plant debris, and the severity of green algae and cyanobacteria at public beaches, lakeside parks, and private piers. Submitted data can be seen in real-time on [Lakeforecast.org](http://Lakeforecast.org) or our free app allowing the general public to stay up to date on current lake conditions.

Data collected from our monitors is used to advocate for implementing watershed improvement projects, raise additional dollars to protect our waters, and prepare our annual [State of the Lakes Report](#) outlining conditions, trends, and water quality drivers affecting the five Yahara lakes.

LakeForecast would not be possible without our dedicated volunteers giving their time and passion to improving Greater Madison’s lakes. Their condition reports highlight the challenges and opportunities when it comes to our lakeside spaces, support important public health initiatives at our public beaches, and improve our understanding of evolving lake dynamics. **Thank you, volunteers!**



**Image 1:** Monitors collect temperature and water clarity readings. **Image 2:** Nearshore training for new 2023 monitors. **Image 3:** Offshore monitors observe zooplankton collected during training with the Center for Limnology.

## 2023 Program Highlights

- 87 nearshore and 5 offshore monitoring stations covering all five Yahara lakes (Figure 1)
- All 25 public beaches monitored at least twice a week
- 90 volunteers contributing 1,910 LakeForecast condition reports
- Algal toxin testing at 10 sites in partnership with UW-Madison Civil & Environmental Engineering
- Partnership with Lake Waubesa Conservation Association to add 4 new monitoring sites
- Partnership with Friends of Lake Kegonsa (FOLKS) to add 3 new monitoring sites
- Volunteer celebration and trivia night at Christy's Landing

## Analysis Summary

During the 2023 monitoring season, low spring and summer rainfall volume and intensity coincided with good water clarity, resulting in higher green algae but fewer cyanobacteria blooms across the five lakes.

This aligns with our understanding of how rainfall drives phosphorus delivery to the lakes, fueling algal blooms. However, precipitation patterns alone are insufficient to explain the conditions seen throughout the spring and summer. Long-term climate trends, the timing of larger rain events, impacts of invasive species, the effect of watershed conservation practices, and spatial variability of monitoring sites make it difficult to identify whole-lake water quality patterns. Watershed improvement projects aimed at reducing phosphorus loading into the lakes often have variable lag times before positive lake responses become noticeable. Clean Lakes Alliance remains dedicated to improving the water quality of the Yahara chain of lakes through partnership building, advocacy efforts, project implementation, and fostering stewardship throughout the watershed.


Please see [Fundamental Concepts on Water Quality of the Yahara Chain of Lakes \(Mendota, Monona, Wingra, Waubesa, and Kegonsa\)](#) for an excellent summary of our current understanding of lake dynamics. These concepts and understandings are also presented in [RENEW THE BLUE: A Community Guide for Cleaner Lakes & Beaches in the Yahara Watershed](#) (2022).



Photo courtesy of monitor Robert Bertera



## 2023 Monitoring Locations



**Table 1:** Nearshore and offshore LakeForecast monitoring effort.

	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
<b>Monitoring Stations</b>	10	46	58	79	69	79	72	71	85	94	87
<b>Volunteers</b>	9	43	57	68	73	87	77	87	101	96	90
<b>Volunteer Hours</b>	224	446	463	610	539	725	544	603	593	579	522
<b>Condition Reports</b>	204	940	1,260	1,698	1,966	2,661	2,045	2,306	2,105	2,094	1,910


## Water Quality Drivers

### Rainfall

Based on weather gauge data from the Dane County Regional Airport, rainfall totals for 2023 (shown in Fig. 2) reflected a very dry start to the summer season. May and June's collective rainfall totaled only 2.01 inches, compared to the normal 9.38 inches. July's 6.21 inches of rain surpassed the monthly normal (4.51 inches), bringing much-needed relief and increased flow through the chain. However, drought conditions remained throughout the summer with August reporting lower-than-normal rainfall (2.42 inches), exasperating already low lake levels.

Additionally, May and June each experienced only one day of rainfall greater than 0.5 inches. Large "gully washing" rain events were minimal this spring and summer, contributing to less early-season runoff. The second half of the summer remained noticeably dry with July and August collectively experiencing only 3 days of rainfall greater than 0.5 inches, with the most notable being August 14th when 2.08 inches of rain was recorded.

This year's continued drought conditions speak to the impacts that runoff has on phosphorus loading and water clarity response. Less stormwater moving across our landscape means less opportunity for polluted runoff to wash into our lakes. It is well documented that early-season rainfall (Jan.-Mar.) typically delivers the most phosphorus runoff to the lakes. As a result, this year's low rainfall totals during that early-season time frame may have helped moderate total phosphorus contributions for the year.



**Figure 2:** Cumulative rainfall collected at the Dane County Regional Airport. Total precipitation collected in inches during May through September compared to historic normals. Monthly precipitation normals are the 30-year average for total monthly precipitation.

## Lake Ice

Winter temperatures were close to normal leading into the 2023 summer monitoring season. Lake Mendota remained frozen for 98 days, or 13 days longer than in 2022. This is on par with the median duration of ice cover on Lake Mendota (102 days) as measured over the last 168 seasons.

A shorter winter leads to warmer water temperatures earlier in the year, effectively creating a longer growing season for algae and aquatic plant life. Warmer winters can also lead to greater runoff and phosphorus delivery when wet precipitation falls on frozen soils and cannot soak into the ground. Rather than infiltrate, rainfall can more easily generate runoff that carries manure and other contaminants toward receiving water bodies. It is estimated that, on average, nearly half of total phosphorus loading through Lake Mendota’s monitored stream tributaries occurs from January to March, making late winter and early spring one of our most vulnerable times for negative water quality impacts.

## Water Clarity

Below-normal rainfall in 2023 meant less runoff washing fertilizers, eroded soil, cow manure, and other sources of phosphorus into our lakes, resulting in comparatively higher clarity conditions. Limited rainfall likely contributed to fewer cyanobacteria blooms than we typically experience aiding high clarity readings.

Table 2 below shows average water clarity recorded on each lake from June 1st - August 31st. **Average nearshore clarity for 2023 was up for all lakes compared to last year’s averages. All lakes except for Mendota reported the highest clarity readings since LakeForecast monitoring started in 2014 (Figure 3). All five lakes fell within the “good” water clarity designation represented in Figure 4.**

**Table 2:** Average nearshore water clarity from 2020 to 2023, collected from June 1st to August 31st as measured with a turbidity tube in centimeters.


	2020			2021			2022			2023		
	June	July	August	June	July	August	June	July	August	June	July	August
<b>Mendota</b>	103	88	81	96	77	65	97	80	75	103	80	75
<b>Monona</b>	102	90	78	89	78	72	108	102	100	107	106	107
<b>Waubesa</b>	110	80	61	112	73	51	110	80	60	106	106	97
<b>Wingra</b>	117	103	104	120	120	86	118	120	120	120	120	120
<b>Kegonsa</b>	90	57	36	94	87	78	100	78	73	116	97	74
<b>All Lakes (Monthly Averages)</b>	104	84	72	102	87	70	107	92	85	110	102	94
<b>All Lakes (Jun-Aug Average)</b>	87			86			95			102		




**Image 4:** The bottom of the lake can be seen far from shore in this photo that showcases 2023’s excellent water clarity. Photo courtesy of monitor Robert Bertera.

# Jun - Aug Average Nearshore Clarity

2014 - 2023



**Figure 3:** Nearshore water clarity from 2014 to 2023 is measured in centimeters using a turbidity tube. The maximum value for the turbidity tube is 120 centimeters, indicating the clearest waters. Lines for average clarity are overlaid in the three different classifications for turbidity: **Good (80-120 cm)**, **Fair (50-80 cm)**, and **Murky (0-50 cm)**.



**Figure 4:** At what depths can you see your toes? We created this adult human representation to help visualize how a lake's water clarity conditions relate to measurements on a turbidity tube.



## Cyanobacteria Blooms & Microcystin Testing

In reviewing this season’s bloom frequency, the same analytical method first adopted in 2018 was used. This approach counts the number of days on each lake with at least one report of a strong cyanobacteria bloom observed within the monitoring sites. By dividing the number of 'cyanobacteria bloom days' by the total number of sampling days for each lake, a percentage is generated representing how often the monitors observed at least one major bloom (Table 3). For example, Lake Kegonsa volunteers reported strong evidence of cyanobacteria on 16% of all monitored days in July 2023. This method lessens overreporting in situations when different monitors report the same cyanobacteria bloom.




Overall, monitors reported **low cyanobacteria blooms during the 2023** sampling period, with most of our significant blooms occurring at the end of May and early June (Figure 5). Monitors on Monona, Waubesa, and Kegonsa did not report a single significant bloom. This is welcomed news after last year’s record-high bloom count on Kegonsa. Lake Wingra experienced similar bloom counts to last year, and Lake Mendota had a handful of significant blooms reported in 2023 after not experiencing any in 2022.

**Image 5:** Cyanobacteria bloom forming at the surface. Photo courtesy of monitor Arlene Koziol.

**Table 3:** Percentage of sampling days with a reported strong presence of cyanobacteria from June 1st to August 31st.

	2020			2021			2022			2023		
	June	July	August	June	July	August	June	July	August	June	July	August
<b>Mendota</b>	10%	3%	0%	13%	26%	6%	0%	0%	0%	3%	0%	0%
<b>Monona</b>	3%	4%	0%	20%	41%	3%	10%	0%	0%	0%	0%	0%
<b>Waubesa</b>	7%	14%	0%	7%	8%	15%	0%	0%	4%	0%	0%	0%
<b>Wingra</b>	0%	0%	0%	0%	0%	0%	0%	6%	0%	0%	0%	6%
<b>Kegonsa</b>	7%	10%	19%	0%	17%	7%	23%	32%	23%	0%	0%	0%
<b>All Lakes (Jun - Aug Average)</b>	6%	7%	4%	9%	20%	7%	8%	8%	6%	1%	0%	1%




**Figure 5:** Percent of sampling days with a strong presence of cyanobacteria for each lake from 2014 to 2023.

These results are consistent with an earlier analysis performed by Dr. Paul Block’s lab at UW-Madison. Dr. Block’s team creates cyanobacteria forecasts for Lake Mendota using geospatial satellite imagery, global weather and ocean circulation patterns, and data collected from our monitors to predict the likelihood of cyanobacteria blooms in our lakes (Figure 6). In early June, Dr. Block’s team estimated that “*despite the early bloom and warm temperatures, our prediction for 2023 reflects a high likelihood of near normal cyanobacteria biomass and a normal number of beach days closed due to elevated cyanobacteria. The models only predict a 15% chance of above-average conditions, largely due to the developing El Niño in the Pacific Ocean*” predicting average cyanobacteria biomass for the season.

A select group of LakeForecast monitors provided supplemental microcystin testing for Dr. Block’s lab at ten locations in 2023. Microcystins are a common class of toxins that can be produced by cyanobacteria, making them unsafe and potentially lethal to people, pets, and wildlife. As a reminder, cyanobacteria (also known as blue-green algae) is not an algae, but a photosynthesizing bacteria naturally occurring in our waters. Only certain strains of cyanobacteria can produce microcystins and species able to generate this toxin do not necessarily produce microcystins during each bloom. In addition, microcystin concentrations are not always correlated with the intensity of the bloom. Having a greater understanding of when cyanobacteria produce microcystins and at what concentration could aid public health officials in assessing toxicity risk.

Monitors reported when blooms were and were not present to randomize collections. Of the 92 samples collected in 2023 across Mendota, Monona, Wingra, and Kegonsa, only 21 tests were positive for the presence of microcystins. Of the tests with positive microcystin results, 10 samples had a microcystin concentration of 10 parts per billion (ppb) or higher. For reference, Public Health of Madison and Dane County close our beaches at 8 ppb or higher.





**Figure 6:** Lake Mendota Cyanobacteria & Beach Closure Experimental Forecast for 2023 calculated from Dr. Paul Block’s lab at UW-Madison.

## Green Algae Blooms

While comparatively fewer cyanobacteria blooms were reported, the same cannot be said for green algae blooms. We had numerous reports of green algae overtaking sampling sites at certain locations. However, the table below shows that **2023 was a typical year regarding historically reported green algae blooms** (Table 4). Lakes Mendota and Monona experienced more significant blooms than last year, while Wingra, Waubesa, and Kegonsa received




fewer reports of blooms compared to 2022 (Figure 7). We utilized the same analytical method for green algae blooms as cyanobacteria blooms. This approach counts the number of days when one or more monitors reported “strong evidence” for a green algae bloom at their monitoring station. By dividing the number of 'green algae bloom days' by the total number of sampling days for each lake, a percentage is generated that better represents the number of blooms witnessed each year on a given lake (Table 4). This method lessens overreporting when different monitors might report the same algal bloom.

**Image 6:** A green algae bloom forming at the surface. Photo courtesy of monitor Sally Lehner.


Increased green algae production is common in high-fertility lakes as clarity increases due to sunlight being able to penetrate deeper into the water column and wider across the lake bed. As shown in Figure 8, there has been a considerable difference in the type of blooms reported by monitors from year to year, with noticeably more green algae blooms than cyanobacteria starting in 2020. While it's perhaps welcomed news to have fewer cyanobacteria blooms across our lakes, green algae blooms are also a symptom of excess phosphorus. At moderate levels, green algae and native aquatic plants play an important role in maintaining healthy lake ecosystems. It's worth noting that green algae on its own is not harmful to human or animal health, but it can be an unwelcome nuisance when this floating organic debris overtakes beaches and piers.

**Table 4:** Rounded percentage of sampling days with a strong presence of green algae reported from June 1st to August 31st.

	2020			2021			2022			2023		
	June	July	August	June	July	August	June	July	August	June	July	August
<b>Mendota</b>	3%	26%	3%	20%	23%	0%	3%	0%	0%	28%	7%	0%
<b>Monona</b>	23%	36%	3%	37%	37%	7%	13%	10%	3%	24%	23%	11%
<b>Waubesa</b>	0%	4%	0%	4%	8%	0%	13%	4%	17%	3%	4%	4%
<b>Wingra</b>	0%	13%	55%	0%	0%	0%	18%	0%	29%	19%	0%	0%
<b>Kegonsa</b>	20%	23%	55%	14%	24%	21%	20%	19%	23%	10%	19%	10%
<b>All Lakes (Monthly Averages)</b>	10%	21%	18%	17%	20%	6%	13%	7%	12%	16%	12%	5%
<b>All Lakes (Jun-Aug Averages)</b>	16%			14%			11%			11%		



**Figure 7:** Percent of sampling days with a strong presence of green algae reported from 2014 to 2023.



**Figure 8:** Percent of sampling days with a strong presence of green algae versus cyanobacteria from 2014 to 2023.

For questions or comments regarding LakeForecast or the material provided, please contact Caitlin McAleavey, Watershed Engagement Manager ([caitlin@cleanlakesalliance.org](mailto:caitlin@cleanlakesalliance.org)), and Paul Dearlove, Deputy Director and Chief Science Officer ([paul@cleanlakesalliance.org](mailto:paul@cleanlakesalliance.org)).

## Thank You, Monitors!

Clean Lakes Alliance wishes to recognize the following volunteer LakeForecast monitors for their many contributions to the success of this program:

### Lake Mendota

Rhonda Arries  
Lynn Ascione  
George Benton  
Mathew Chotlos  
Emily Conklin  
Bennett Davishoff  
Doris Dubielzig  
Jan Eddy  
Nancy Ellison  
Friends of UW Lakeshore  
Nature Reserve

Robert Gilbert  
Bob Gross  
Mary Hillebrand  
Andy Ippolito  
Jerry Jendrisak  
Sydney Kerstein  
Arlene and Jeff Koziol  
Jackie Lanear  
Jay Larson  
Peggy Lee  
Allan Levin  
Linda Malkin

Seth McGee  
Carol Michalski  
MJ and Tom Morgan  
Stephanie Ogborne  
Raymond Rajala  
Cole Seckel  
Kim Sprecher  
Jacob Stampen  
Mattie Urrutia  
Will Vuyk  
Angela Webster  
Ian Wegger



**Lake Monona**

Allen Arntsen  
Doug Bach  
Jill Ball  
Bob Bertera  
Carolyn Betz  
Brittingham Boats  
Claire Finucane  
Amie Francisco  
Rachel Groman  
Joanne Kaminsky  
Renee Long  
Mel McCartney  
Elaine Meier  
Joy Newmann  
Austin Novak  
Michael Oleniczak  
Reed Rodenkirch  
Zak Redding  
David Schroder  
Tom Smith  
Christine Stocke  
Jan Van Vleck  
Sam Warfel  
Amy Wencel  
Paul Zech

Tam Knickmeier  
John McIntyre  
Mike Moran  
Tom Reuschlein  
Lori Wallman  
Kimberly Walsh

**Lake Kegonsa**

Sarah Balz  
Bill & Amanda Blank  
John Bottorff  
George Corrigan  
Connie Hagen  
Janice Kellogg  
Ken LePine  
Anne Meisner  
Terry Peterson  
Claudia and Greg Quam  
Gary Tapper  
Peg Stine and Carl Johnson  
Kim Van Brocklin  
Kim & Tom Walz  
Jim Wilcox

**Lake Wingra**

Regan Botsford  
Lisa Grueneberg  
Jess Hankey  
Cindy Koschmann  
Sally Lehner  
John Nicol

**Lake Waubesa**

Kitty Brussock  
Cassy Cooley  
Troy Hibicki  
Mike Horn  
Marissa Hueffner  
Ginny Kominiak  
Woody Kneppreth

